KNOW YOUR CONCUSSION ABB CS Be alert for Contact a

Assess the situation

Be alert for Co signs and he symptoms pr

for Contact a d health care ns provider

What is a concussion?

A concussion is a type of brain injury that changes

the way the brain normally works. A concussion is

be a mild bump to the head can be serious.

caused by a bump, blow, or jolt to the head and can also

occur from a blow to the body that causes the head and

brain to move rapidly back and forth. Even what seems to

Concussions can have a more serious effect on a young,

developing brain and need to be addressed correctly.

Wisconsin Fact Sheet for Parents

What are the signs and symptoms of a

concussion?

You can't see a concussion. Signs and symptoms of concussion can show up right after an injury or may not appear or be noticed until hours or days after the injury. It is important to watch for changes in how your child or teen is acting or feeling, if symptoms are getting worse, or if s/he just "doesn't feel right." Most concussions occur without loss of consciousness.

If your child or teen reports **one or more** of the symptoms of concussion listed below, or if you notice the symptoms yourself, seek medical attention right away. Children and teens are among those at greatest risk for concussion.

GNS OBSERVED BY PARENTS OR GURADIA	NS SYMPTOMS REPORTED BY YOU	R CHILD
Appears dazed or stunned Is confused about events Answers questions slowly Repeats questions Can't recall events prior to the hit, bump, or fall Can't recall events after the hit, bump, or fall Loses consciousness (even briefly) Shows behavior or personality changes Forgets class schedule or assignments	 Thinking/Remembering: Difficulty thinking clearly Difficulty concentrating or remembering Feeling more slowed down Feeling sluggish, hazy, foggy, or groggy Physical: Headache or "pressure" in head Nausea or vomiting Balance problems or dizziness Fatigue or feeling tired Blurry or double vision Sensitivity to light or noise Numbness or tingling Does not "feel right" 	Emotional: Irritable Sad More emotional than usual Nervous Sleep*: Drowsy Sleeps less than usual Sleeps more than usual Has trouble falling asleep *Only ask about sleep symptoms if the injury occurred on a prior day.

PUBLIC INSTRUCTION



DANGER SIGNS

Be alert for symptoms that worsen over time. Your child or teen should be seen in an emergency department right away if s/he has:

- One pupil (the black part in the middle of the eye) larger than the other
- Difficult to arouse
- Severe headache or worsening headache
- · Weakness, numbness, or decreased coordination
- Repeated vomiting or nausea
- Slurred speech
- Convulsions or seizures
- Difficulty recognizing people or places
- · Increasing confusion, restlessness, or agitation
- Unusual behavior
- Loss of consciousness (even a brief loss of consciousness should be taken seriously)



Children and teens with a suspected concussion should NEVER return to sports or recreation activities on the same day the injury occurred. They should delay returning to their activities until a health care provider experienced in evaluating for concussion says they are symptom-free and provide written clearance to return to activity. This means, until permitted, not returning to:

- Physical Education (PE) class,
- Sports conditioning, weight
- lifting, practices and games, or
- Physical activity at recess.

What should I do if my child or teen has a concussion?

- Seek medical attention right away. A health care provider experienced in evaluating for concussions can direct concussion management and review when it is safe for your child to return to normal activities, including school (concentration and learning) and physical activity. If your child or teen has been removed from a youth athletic activity because of a suspected or confirmed concussion or head injury, they may not participate again until he/she is evaluated by a health care provider and receives written clearance to participate in the activity from the health care provider.
- Help them take time to get better. If your child or teen has a concussion, her or his brain needs time to heal. Your child or teen should limit activities while he/she is recovering from a concussion. Exercising or activities that involve a lot of concentration, such as studying, using a computer, texting, or playing video games may worsen or prolong concussion symptoms (such as headache or tiredness). Rest will help your child recover more quickly. Your child may become upset that he/she cannot participate in activities.

 Together with your child or teen, learn more about <u>concussions</u>. Talk about the potential long-term effects of concussion and the problems caused by returning too soon to daily activities to quickly (especially physical activity and learning/concentration).

How can I help my child return to school safely after a concussion?

Help your child or teen get needed support when returning to school after a concussion. Talk with your child's school administrators, teachers, school nurse, coach, and counselor about your child's concussion and symptoms. Your child may feel frustrated, sad, and even angry because he/she cannot keep up with schoolwork and learn as well after a concussion. Your child may also feel isolated from peers and social networks. Talk often with your child about these issues and offer your support and encouragement. As your child's symptoms decrease, the extra help or support can be removed gradually. Children and teens who return to school after a concussion may need to:

- Take rest breaks as needed,
- Spend fewer hours at school,
- Be given more time to take tests or complete assignments,
- Receive help with schoolwork, and/or
- Reduce time spent reading, writing, or on the computer.

To download this fact sheet in Spanish, please visit: www.cdc.gov/Concussion. Para obtener una copia electrónica de esta hoja de información en español, por favor visite: www.cdc.gov/Concussion.

www.cdc.gov/Concussion; www.wiaawi.org; www.nfhs.org